

ON THE NON-ABELIAN BORN-INFELD ACTION

Pascal Bain

*Laboratoire de Physique Théorique de l'École Normale Supérieure[†]
24 rue Lhomond, F-75231 Paris Cedex 65.*

Abstract We discuss some aspects of the generalization of the Born-Infeld action to non-abelian gauge groups and show how the discrepancy between Tseytlin's symmetrized trace proposal and string theory can be corrected at order F^6 . We also comment on the possible quadratic order fermionic terms.

While the talk I gave at the Gong Show session of Cargèse' 99 Summer School was devoted to an outline of the main results of the paper "Curvature terms in D-brane actions and their M-theory origin" [1], this note will be focused on another very interesting but not yet settled aspect of the D-brane effective action : the generalization of the Dirac-Born-Infeld (DBI) action to non-abelian gauge groups.

The massless degrees of freedom of an isolated D-brane are its transverse coordinates, a world-volume $U(1)$ gauge field, and the associated fermions [2]. Its effective action contains a Dirac-Born-Infeld (DBI) (see [3] for a recent review of the DBI action) and a Wess-Zumino (WZ) piece. While the WZ part, describing the interactions of the D-brane with Ramond-Ramond field backgrounds, can be determined by anomaly cancellation and is therefore believed to be exact, the DBI piece is only reliable for small space-time curvatures, and in the low-acceleration regime. Indeed, for non constant gauge field strength, the DBI action is corrected by derivative terms [3, 4].

When N identical D-branes coincide, the transverse coordinates and world-volume gauge fields become $N \times N$ unitary matrices. The non-abelian generalization of the DBI action describing this dynamics has

*LPTENS-99/34

[†]UMR 8549 du Centre National de la Recherche Scientifique et de l'École Normale Supérieure.

been the subject of several recent papers. In the low-energy non-relativistic limit one must recover the usual quadratic action of dimensionally-reduced super Yang-Mills. The question is whether there exist less-trivial truncations, taking into account relativistic effects in a controlled manner. However, since $[\mathbf{D}_m, \mathbf{D}_n]\mathbf{F}_{pq} = [\mathbf{F}_{mn}, \mathbf{F}_{pq}]$ for a non-abelian group, the part of the action which depends on the field strength but not on its covariant derivative is ambiguous, contrary to the abelian case. One simple proposal for the flat space-time action has been put forward by Tseytlin [5] :

$$\mathcal{L} = \mathbf{STr} \sqrt{-\det(\eta + 2\pi\alpha'\mathbf{F})} \quad (1.1)$$

The prescription is to expand the Born-Infeld action as a formal power series, symmetrize under all permutations of the (non-abelian) field strengths and then take the trace. The resulting action captures several features of the full string theory.

First, since $\mathbf{STr}(t_8\mathbf{F}^4) = \mathbf{Tr}(t_8\mathbf{F}^4)$, it reproduces the already well-known F^4 which can be obtained from a direct four-point function on the disk diagram [6] or by a two-loop β -function for the non-abelian background gauge field in open string theory [7].

Second, in four dimensions, it reduces to the quadratic Yang-Mills term for an (anti-)self-dual field strength as pointed out in [8] where BPS states of the non-abelian BI action were investigated.

Since it seems out of reach to test this proposal to higher orders in α' by a direct string theory computation – it would involve a six-point open string amplitude on the disk or a four-loop β function! – one must use indirect arguments. Hashimoto and Taylor have found a configuration [9] which exhibits disagreements with the symmetrized trace prescription : they considered a constant magnetic background field [10] or, in the dual string picture, a pair of intersecting rotated D2-branes living on a torus. To the background considered in [9], it is possible to add a $U(1)$ part which corresponds to the simultaneous rotation of the two branes in the dual picture. We expect that the spectrum still only depends on the relative orientation of the branes. This abelian gauge field imposes further constraints on the form of the action.

For an (anti-)self-dual background, the spectrum obtained with the symmetrized trace prescription exactly matches the string theory spectrum, a result not fully realized in [9]. For more general constant fields, both results agree up to order $(\alpha')^4$, which is consistent with the fact that the symmetrized trace is the correct answer for F^4 terms, as already pointed out. The discrepancy appears at the sixth order and can be solved at this order by adding terms with commutators of the fields strength $\mathbf{F} = F^a\mathbf{T}^a$ to the action (1.1). For simplicity, we have only

considered terms with two commutators, but even with this restriction, the test does not completely fix the action. If we define the tensors

$$\begin{aligned} C^{\pm a_1 a_2 a_3 a_4 a_5 a_6} &= \text{Tr}(\mathbf{T}^{a_1} \mathbf{T}^{a_2} \{[\mathbf{T}^{a_3}, \mathbf{T}^{a_4}], [\mathbf{T}^{a_5}, \mathbf{T}^{a_6}]\})/4 \\ &\pm \text{Tr}(\mathbf{T}^{\{a_1, [\mathbf{T}^{a_3}, \mathbf{T}^{a_4}] \mathbf{T}^{a_2}\}} [\mathbf{T}^{a_5}, \mathbf{T}^{a_6}])/4 \end{aligned}$$

and the contractions

$$\begin{aligned} V_1^\pm &= (2\pi\alpha')^6 F_{mn}^{a_1} F_{np}^{a_2} F_{pq}^{a_3} F_{rm}^{a_4} F_{qs}^{a_5} F_{sr}^{a_6} C^{\pm a_1 a_2 a_3 a_4 a_5 a_6} \\ V_2^\pm &= (2\pi\alpha')^6 F_{mn}^{a_1} F_{pq}^{a_2} F_{nr}^{a_3} F_{sm}^{a_4} F_{rp}^{a_5} F_{qs}^{a_6} C^{\pm a_1 a_2 a_3 a_4 a_5 a_6} \\ V_3^\pm &= (2\pi\alpha')^6 F_{mn}^{a_1} F_{pq}^{a_2} F_{np}^{a_3} F_{rs}^{a_4} F_{qm}^{a_5} F_{sr}^{a_6} C^{\pm a_1 a_2 a_3 a_4 a_5 a_6} \\ V_4^\pm &= (2\pi\alpha')^6 F_{mn}^{a_1} F_{pq}^{a_2} F_{nm}^{a_3} F_{qr}^{a_4} F_{sp}^{a_5} F_{rs}^{a_6} C^{\pm a_1 a_2 a_3 a_4 a_5 a_6} \\ V_5^\pm &= (2\pi\alpha')^6 F_{mn}^{a_1} F_{pq}^{a_2} F_{nm}^{a_3} F_{rs}^{a_4} F_{qp}^{a_5} F_{sr}^{a_6} C^{\pm a_1 a_2 a_3 a_4 a_5 a_6} \end{aligned}$$

the string theory spectrum is reproduced at order six in α' when we add

$$\frac{1}{24}(V_1^+ - V_2^+ - \frac{1}{2}V_3^+ - V_4^+ + \frac{1}{4}V_5^+) + \frac{1}{360}(V_1^- - V_2^- - \frac{1}{2}V_3^- - V_4^- + \frac{1}{4}V_5^-)$$

to the action (1.1). We have also checked that for an (anti-)self-dual field strength, these additional terms vanish and, then, the effective action reduces to the linear Yang-Mills action.

This situation seems somewhat similar to the problem studied by Douglas, Kato and Ooguri [11] where they demonstrate the necessity to add commutators terms to the symmetrized trace in their D-Kähler potential (at sixth order) to reproduce the mass-shell conditions.

A more exhaustive and systematic investigation, combining the self-duality and Hashimoto-Taylor's constraints, would be desirable. For this purpose, it seems convenient to use the following diagrammatic representation to enumerate all the possible terms appearing at a given order $(\alpha')^n$: the idea is to put each $\mathbf{F}$ field at a vertex of a polygon of degree n and to symbolize each Lorentz contraction by a line joining these points. The trace over the gauge group is taken in the standard order on the polygon. For example, the $\mathbf{STr}(\mathbf{F}^4)$ terms are given in figure (1.1).




Figure 1.1 Diagrammatic representation of $3 \mathbf{STr}(\mathbf{F}^4)$.




Figure 1.2 Relation between diagrams for $F = *F$.

Notice that in the symmetrized trace, the weight of each term for a given set of Lorentz contractions are just the number of non-superimposable diagrams under rotations preserving the polygon vertices : in this way, in figure (1.1), the first and third diagrams have weight two and the second and fourth weight one. Using this representation, one can easily enumerate the possible terms at order $(\alpha')^6$; there are exactly twenty eight inequivalent diagrams : five of the “ $(F^2)^3$ ” type, nine “ $(F^2)(F^4)$ ” and fourteen “ F^6 ”. Moreover, for a self-dual field strength, we have the relation :

$$2(\mathbf{F}_{mp}\mathbf{F}_{pn} + \mathbf{F}_{np}\mathbf{F}_{pm}) = \eta_{mn}\mathbf{F}_{pq}\mathbf{F}_{qp}$$

which can be translated into the diagrammatic equation of figure (1.2). Since for a self-dual field strength higher-order terms should vanish [8], this implies further constraints on the possible form of the action. However, using this rule, it is quite simple to analyse these constraints; for example, it is easy to verify that the F^4 term in figure (1.1) vanishes for a self-dual F .

Let us conclude with some remarks : first, the effective action should also contain fermionic terms. Aganagic, Popescu and Schwarz have proposed a supersymmetric generalization of the abelian BI action [12]. To study its generalization to non-abelian groups, we can use a test analogous to the one developed for the gauge field in [9]. From the dual string theory, one expect, for fermions in a constant gauge field, a Landau spectrum of the form :

$$2\Sigma_{12} \cdot \arctan(2\pi\alpha' F_{12}) + 2\Sigma_{34} \cdot \arctan(2\pi\alpha' F_{34})$$

with $\Sigma_{ij} = [\Gamma_i, \Gamma_j]$. Such spectrum can be obtained from the Lagrangian :

$$\mathcal{L}_2 = \mathbf{STr} \left[\overline{\lambda'} (\eta + 2\pi\alpha' \mathbf{F})_S^{-1\mu\nu} \Gamma_\mu \mathbf{D}_\nu \lambda' \right].$$

Finally, Seiberg and Witten [13] have recently proved that the standard abelian BI action is equivalent, up to terms involving derivatives of F , to the abelian BI for a non-commutative gauge field by a field

redefinition. This result should still hold for the non-abelian theory. It would be interesting to see what the symmetrized trace prescription on one side gives on the other side. Moreover, in the limit $\alpha' \rightarrow 0$ considered in [13], the action on the non-commutative side reduces to the non-commutative Yang-Mills Lagrangian whereas it remains non-trivial on the commutative side. Since the generalization of this abelian NCYM to the non-abelian case is trivial, one can ask how this discussion can be extended and give some insights on the commutative non-abelian side.

Acknowledgments

I would like to thank the organizers of the Cargèse'99 ASI for a very nice school and for financial support. It is a pleasure to thank C. Bachas for collaboration on this subject.

References

- [1] C.P. Bachas, P. Bain, M.B. Green, JHEP **9905** (1999) 011.
- [2] J. Polchinski, *TASI Lectures on D-Branes* [hep-th/9611050]; C.Bachas, *Lectures on D-branes* [hep-th/9806199].
- [3] A.A. Tseytlin, “Born-Infeld action, supersymmetry and string theory”, [hep-th/9908105].
- [4] O.D. Andreev, A.A. Tseytlin, Nucl.Phys. **B311** (1988) 205.
- [5] A.A. Tseytlin, Nucl.Phys. **B501** (1997) 41
- [6] D.J. Gross and E. Witten, Nucl.Phys. **B277** (1986) 1.
A. A. Tseytlin, Nucl.Phys. **B276** (1986) 391; Nucl.Phys. **B291** (1987) 876 (E).
- [7] D. Brecher, M.J. Perry, Nucl.Phys. **B527** (1998) 121.
- [8] D. Brecher, Phys.Lett. **B442** (1998) 117.
- [9] A. Hashimoto, W. Taylor IV, Nucl.Phys. **B503** (1997) 193
- [10] A. Abouelsaood, C. Callan, C. Nappi, S. Yost, Nucl.Phys. **B280** (1987) 123.
- [11] M.R. Douglas, A. Kato, H. Ooguri, [hep-th/9708012].
- [12] M. Aganagic, C. Popescu, J.H. Schwarz, Nucl.Phys. **B495** (1997) 99.
- [13] N. Seiberg, E. Witten, [hep-th/9908142].